

Handgrip Strength and Serum Uric Acid Levels are Associated with Dialysis-Malnutrition Scores in Chronic Hemodialysis Patients

El-Sıkma Gücü ve Serum Ürik Asit Seviyeleri, Kronik Hemodiyaliz Hastalarında Diyaliz-Malnütrisyon Skoru ile İlişkilidir

ABSTRACT

OBJECTIVE: Protein-energy-wasting (PEW) is common and contributes to increased mortality in hemodialysis (HD) patients. Herein we aimed to investigate the relationships between the parameters of malnutrition including dialysis malnutrition score (DMS), bioimpedance analysis (BIA), handgrip strength (HGS) in chronic HD patients.

MATERIAL and METHODS: Forty chronic HD patients (mean age:58±18 years; males:22) were enrolled. Mid-arm circumference (MAC) measurements and BIA were performed. HGS was measured by a Jamar hydraulic hand dynamometer.

RESULTS: HGS, MAC and uric acid levels were higher in patients with mild malnutrition compared to severe malnutrition group. Postdialysis creatinine was associated with lean body ratio (LBR) ($r=0.56$, $p=0.002$), DMS ($r=-0.57$, $p<0.001$), HGS ($r=0.73$, $p<0.001$) and MAC ($r=0.51$, $p=0.001$). DMS was inversely correlated with HGS ($r=-0.63$, $p<0.001$), MAC ($r=-0.48$, $p=0.002$), total protein ($r=-0.36$, $p=0.02$) and uric acid levels ($r=-0.54$, $p<0.001$). In linear regression analysis ($R^2=0.58$; $p<0.001$), HGS and uric acid were found to independently predict DMS. HGS was associated with MAC, LBR, uric acid and post-HD urea. HGS was negatively correlated with age, DMS, fat tissue ratio (FTR) and Kt/V. In linear regression analysis ($R^2=0.71$; $p<0.001$), gender, DMS and MAC were found to independently predict HGS.

CONCLUSION: HGS and uric acid were significantly associated with dialysis-malnutrition scores. HGS is an easy and reliable test for the evaluation of malnutrition in HD patients.

KEY WORDS: Malnutrition, Hemodialysis, Handgrip strength, Dialysis malnutrition score, Uric acid

ÖZ

AMAÇ: Protein-enerji tükenmesi sendromu, hemodiyaliz (HD) hastalarında sıklık ve artmış mortalite ile ilişkilidir. Burada, el-sıkma gücünün (ESG); diyaliz malnütrisyon skoru (DMS) ve biyoimpedans analizi (BIA) gibi diğer malnütrisyon parametreleri ile ilişkisinin araştırılması amaçlanmıştır.

GEREÇ ve YÖNTEMLER: Kırk kronik HD hastası (ortalama yaş: 58±18 yıl; erkek :22) çalışmaya katıldı. Orta-kol çevresi (OKÇ) ve BIA ölçümleri yapıldı. ESG, Jamar hidrolik el dinamometresi ile ölçüldü.

BULGULAR: ESG, OKÇ ve ürik asit seviyeleri, hafif malnütrisyon grubunda, ciddi malnütrisyon grubuna göre daha yüksek saptandı. Postdiyaliz kreatinin seviyeleri, yağsız vücut oranı (YVO) ($r=0.56$, $p=0.002$), DMS($r=-0.57$, $p<0.001$), ESG($r=0.73$, $p<0.001$) ve OKÇ($r=0.51$, $p=0.001$) ile anlamlı olarak ilişkili bulundu. DMS ise ESG ($r=-0.63$, $p<0.001$), OKÇ($r=-0.48$, $p=0.002$), total protein($r=-0.36$, $p=0.02$) ve ürik asit seviyeleri ($r=-0.54$, $p<0.001$) ile ters-ilişkili bulundu. Lineer regresyon analizinde ($R^2=0.58$; $p<0.001$), ESG ve ürik asitin DMS'nin bağımsız öngörütücüleri olduğu bulundu. ESG ile OKÇ, YVO, ürik asit ve post-HD üre değerleri ilişkiliydi. Buna karşın ESG; yaş, DMS, yağ doku oranı ve Kt/V ile ters ilişkili bulundu. Lineer regresyon analizinde ($R^2=0.71$; $p<0.001$), cinsiyet, DMS ve OKÇ'nin ESG'nin bağımsız öngörütücüleri oldukları gösterildi.

SONUÇ: ESG ve ürik asit seviyeleri, diyaliz-malnütrisyon skorlarıyla anlamlı olarak ilişkilidir. ESG, HD hastalarında malnütrisyonun değerlendirilmesi için kolay ve güvenilir bir test olabilir.

ANAHTAR SÖZCÜKLER: Malnütrisyon, Hemodiyaliz, El sıkma gücü, Diyaliz malnütrisyon skoru, Ürik asit

Semih BAŞCI
Abdullah ÖZKÖK
Mevlüt KAÇAR
Satılmış BİLGİN
Ali Rıza ODABAŞ

Istanbul Medeniyet University,
Goztepe Training and Research Hospital,
Department of Internal Medicine and
Nephrology,
Istanbul, Turkey



Received : 10.03.2017

Accepted : 19.06.2017

Correspondence Address:

Abdullah ÖZKÖK
Istanbul Medeniyet Üniversitesi,
Göztepe Eğitim ve Araştırma Hastanesi
İç Hastalıkları, Nefroloji Bölümü,
Istanbul, Turkey
Phone : + 90 530 828 64 12
E-mail : abduallahozkok@yahoo.com

INTRODUCTION

Protein-energy wasting (PEW) is described as the reduction in protein and energy reserves and it contributes to extremely increased morbidity and mortality in patients with CKD (1). PEW is commonly observed in CKD patients, and its prevalence ranges from 23% to 76% (2). However, the diagnosis of PEW may be difficult. Various clinical, anthropometric and biochemical parameters have been determined as the criteria of PEW such as nutritional scoring systems (such as dialysis malnutrition score- DMS), low body mass index (BMI), muscle mass, decreased serum albumin, and transthyretin and cholesterol levels (1). In CKD patients, measurements of BMI are commonly interfered with by hypervolemia and thus determination of lean body mass and muscle mass is not easy. Body composition measurements using bioimpedance analysis (BIA) are therefore preferred to evaluate loss of lean body and muscle mass, which are considered reliable indicators of PEW (3). Besides these parameters, functional evaluation of the muscle mass is also important. The handgrip strength test (HGS) which is an indicator of the maximal voluntary force of the hand, is a noninvasive, objective and rapid test to evaluate muscle function (4). HGS has been found to reflect the nutritional status of various patient groups including CKD patients (5-7). Herein we aimed to investigate the relationships between the various parameters of malnutrition including DMS, body composition measurements with BIA, functional evaluation of muscle mass with HGS and adequacy of dialysis in chronic hemodialysis (HD) patients.

MATERIALS and METHODS

Study Population and Data Collection

A total of 40 chronic HD patients (mean age: 58 ± 18 years; male:22/female:18) were enrolled. Median time (IQR) on dialysis was 82(22-135) months. Etiologies of chronic kidney disease were as follows: diabetes mellitus, 7 (17.5%); hypertension, 7 (17.5%); glomerulonephritis, 5 (12.5%); polycystic kidney disease, 3 (7.5%); others, 5 (12.5%) and unknown, 13 (32.5%). Patients were receiving thrice weekly dialysis for 4-hour periods with a single-use, hollow-fiber, high-flux HD membrane (Nipro-ELISIO, Osaka, Japan). Blood flow rates ranged from 250 to 300 mL/min, while the dialysate flow rate was kept constant at 500 mL/min. The standard dialysate solution was composed of glucose 1 g/L, bicarbonate: 32 mmol/L, sodium 140 mmol/L, potassium 2 mmol/L; chloride: 111 mmol/L, calcium 1.25 mmol/L and magnesium 0.5 mmol/L. Mean eKt/V was found to be 1.61 ± 0.47 . Mean urea reduction rate (URR) was $75 \pm 10\%$.

The exclusion criteria were determined as follows; age older than 90 and younger than 18 years, presence of pregnancy, active malignancy, infection, advanced heart failure, inflammatory diseases including inflammatory bowel disease. Also patients with cardiac pacemaker and a history of amputation were excluded.

This study was reviewed and approved by the local ethical committee (No: 2015/0149). This study was performed according to good medical and laboratory practices and the recommendations of the Declaration of Helsinki on Biomedical Research Involving Human Subjects. Informed consents were obtained from all participants of the study.

Laboratory Analysis

Fasting blood samples were collected before the midweek HD session. Laboratory values such as complete blood cell counts, electrolytes, uric acid, calcium, phosphorus, total protein, albumin, total cholesterol, triglycerides, iron, ferritin, C-reactive protein and intact parathyroid hormone (PTH) were measured. Serum levels of urea and creatinine were measured before and after dialysis. Accordingly urea reduction rate and equilibrated Kt/V parameters were calculated.

Anthropometric Measurements

Body mass index was calculated as weight/ (height)² of the patient. Mid-arm circumference (MAC) was measured. HGS was measured at least 3 times by a Jamar hydraulic hand dynamometer (Patterson Medical, IL, US) with the shoulder adducted and elbow flexed at 90° and the highest value was accepted as the HGS of the patient.

Bioelectrical Impedance Analysis (BIA) Measurements

Bioelectrical impedance analysis was performed to determine the body composition including fat tissue ratio (FTR) and lean body ratio (LBR). FTR and LBR measurements were done after the HD sessions during the “dry-weight” period to avoid the possible confounding effects of inter-variability of volume status between the patients in the pre-dialysis period.

Dialysis Malnutrition Score

The malnutrition degree was evaluated with subjective global assessment (SGA) as described by Kalantar-Zadeh et al. (8). Accordingly, DMS consists of seven features: weight change, dietary intake, gastrointestinal symptoms, functional capacity, comorbidity, subcutaneous fat and signs of muscle wasting. Each component has a score from 1 (no change) to 5 (very severe malnutrition). The DMS is defined as a number between 7 (normal) and 35 (severely malnourished). Patients were divided into two groups according to DMS such as the no malnutrition to mild malnutrition group (DMS: 7-13) and the moderate to severe malnutrition group (14-35).

Statistical Analysis

Statistical analysis was performed with the Statistical Package for Social Sciences for Windows version 16.0 (SPSS Inc, Chicago, IL, USA). Results of the data with normal distribution were expressed as mean \pm standard deviation. Data with non-normal distribution were presented as median (interquartile range: 25-75%). Comparisons of the groups were performed using the t test or Mann-Whitney U test. Correlation

analysis was performed by Pearson's test and Spearman's test where appropriate. Linear regression analysis was performed to evaluate the variables predicting DMS and HGS. Two-tailed p value of <0.05 was defined as statistically significant.

RESULTS

Biochemical and malnutrition parameters of the patients are presented in Table I. Comparison of male and female patients in terms of biochemical and malnutrition parameters is presented in Table II. Accordingly, HGS (Figure 1), LBR and serum uric acid levels were significantly higher in males compared to female patients. Conversely, FTR, Kt/V, URR, total cholesterol and HDL levels were found to be significantly higher in females. Age and DMS tended to be higher in female patients (p=0.09 and p=0.06 respectively). BMI and MAC were found to be similar in male and female patients.

Patients were divided into two groups according to DMS scores such as the no malnutrition to mild malnutrition group (DMS: 7-13) and the moderate to severe malnutrition group (DMS: 14-35). Comparison of the groups is presented in Table III. BMI, HGS, MAC and serum uric acid levels were found to be significantly higher in patients with mild malnutrition compared to severe malnutrition. On the other hand, patients were older and they had higher Kt/V and URR values in the severe malnutrition group compared to the mild malnutrition group.

Predialysis serum creatinine levels were significantly associated with LBR (r=0.48, p=0.008), DMS (r= -0.39, p=0.01), HGS (r=0.44, p=0.005), hemoglobin (r=0.45, p=0.004), uric acid (r=0.46, p=0.003), total protein (r=0.40, p=0.01) and




Figure 1: Handgrip strength was significantly higher in male patients compared to female patients (p<0.001).

Table I: Biochemical and malnutrition parameters of the patients.

Parameters	Mean (±SD) / median (IQR)
Age (years)	58 ± 18
Dx vintage (months)	82 (22-135)
BMI (kg/m ²)	24 ± 5
DMS	13 ± 4
MAC (cm)	26.75 ± 4.38
Handgrip strength (kg)	22.6 ± 11.34
Lean body ratio (%)	37 ± 9
Fat tissue ratio (%)	27 ± 13
Total leukocyte (/mm ³)	6692 ± 1709
Lymphocyte count (/mm ³)	1626 ± 486
Hemoglobin (g/dl)	11.1 ± 1.36
Iron (µg/dL)	62.98 ± 20.31
Ferritin (ng/mL)	554 ± 281
Parathormone (pg/mL)	606 (386-771)
CRP (mg/dL)	0.95 (0.40-1.67)
Kt/V (equilibrated)	1.61 ± 0.47
URR (%)	75 ± 10
Urea (Pre-dx) (mg/dL)	131 ± 39
Urea (Post-dx) (mg/dL)	34 ± 17
Creatinine (Pre-dx) (mg/dL)	8.98 ± 2.84
Creatinine (Post-dx) (mg/dL)	3.14 ± 1.47
Uric acid (mg/dL)	5.93 ± 1.38
Calcium (mg/dL)	8.38 ± 1.09
Phosphorus (mg/dL)	5.48 ± 1.31
Total Protein (g/dL)	6.61 ± 0.59
Albumin (g/dL)	3.58 ± 0.32
Total cholesterol (mg/dL)	168 ± 39
Triglyceride (mg/dL)	144 ± 64
LDL (mg/dL)	103 ± 31
HDL (mg/dL)	36 ± 12

Dx: Dialysis, **BMI:** Body mass index, **DMS:** Dialysis malnutrition score, **MAC:** Mid-arm circumference, **CRP:** C-reactive protein, **URR:** Urea reduction rate, **LDL:** Low-density lipoprotein, **HDL:** High-density lipoprotein

albumin ($r=0.44$, $p=0.005$). However, they were not associated with BMI ($r=0.16$, $p=0.32$). Postdialysis serum creatinine levels were significantly associated with age ($r= -0.38$, $p=0.01$), LBR ($r=0.56$, $p=0.002$) (Figure 2), DMS ($r= -0.57$, $p<0.001$), HGS ($r=0.73$, $p<0.001$) (Figure 3), MAC ($r= 0.51$, $p=0.001$), hemoglobin ($r=0.35$, $p=0.03$) and uric acid ($r=0.57$, $p<0.001$).

Postdialysis urea levels were associated with DMS ($r= -0.37$, $p=0.01$), MAC ($r=0.49$, $p=0.001$), HGS ($r=0.51$, $p=0.001$), and BMI ($r=0.37$, $p=0.02$) but predialysis urea levels were not related to the mentioned parameters.

Correlation analysis between DMS and various anthropometric and biochemical parameters is presented in Table IV. Accordingly, DMS was significantly associated with age ($r=0.43$, $p=0.005$) and Kt/V ($r=0.39$, $p=0.01$). In contrast, DMS was inversely correlated with HGS ($r= -0.63$, $p<0.001$) (Figure 4), MAC ($r= -0.48$, $p=0.002$), serum total protein ($r= -0.36$, $p=0.02$) and uric acid levels ($r= -0.54$, $p<0.001$) (Figure 5).

Linear regression analysis (Adjusted $R^2 = 0.58$; $p<0.001$) was performed to determine factors independently predicting DMS (Table V). Age, gender, HGS, MAC, LBR, total protein, uric acid and Kt/V were included in the analysis as variables. HGS and serum uric acid levels were found to be the independent variables significantly predicting DMS.

HGS was significantly associated with MAC, LBR (Figure 6), serum uric acid and post HD serum urea levels. It tended to be related to serum phosphorus levels ($p=0.05$). In contrast, HGS was negatively correlated with age, DMS, FTR, Kt/V, URR and serum HDL concentrations (Table VI).

Linear regression analysis (Adjusted $R^2 = 0.71$; $p<0.001$) was performed to determine the factors independently predicting HGS (Table VII). Age, gender, DMS, MAC, LBR, FTR, Kt/V, uric acid, HDL and ferritin were included in the analysis. Gender, DMS and MAC were found to be the independent variables significantly predicting HGS.

Table II: Comparison of male and female patients in terms of biochemical and malnutrition parameters.

	Male (n=22)	Female (n=18)	p
Age (years)	53 ± 17	63 ± 18	0.09
Dx vintage (months)	83 (41-141)	60 (20-155)	0.73
BMI (kg/m ²)	24 ± 5	24 ± 6	0.89
DMS	12 ± 4	15 ± 4	0.06
Handgrip strength (kg)	28.36 ± 10.95	15.55 ± 7.18	<0.001
MAC (cm)	27.09 ± 3.95	26.33 ± 4.95	0.59
Lean body ratio (%)	42 ± 7	31 ± 7	<0.01
Fat tissue ratio (%)	20 ± 10	34 ± 11	0.002
KT/V	1.42 ± 0.32	1.84 ± 0.53	0.007
URR (%)	71 ± 8	79 ± 11	0.02
Lymphocyte count (/mm ³)	1648 ± 414	1601 ± 575	0.77
Urea (Pre-dx) (mg/dL)	139 ± 46	120 ± 25	0.10
Creatinine (Pre-dx) (mg/dL)	10.43 ± 1.94	7.18 ± 2.78	<0.001
Hemoglobin (g/dL)	11.41 ± 1.39	10.75 ± 1.27	0.12
Uric acid (mg/dL)	6.34 ± 1.17	5.42 ± 1.48	0.03
Albumin (g/dL)	3.61 ± 0.27	3.52 ± 0.38	0.36
Total cholesterol (mg/dL)	157 ± 30	182 ± 46	0.04
Triglyceride (mg/dL)	146 ± 66	142 ± 64	0.84
LDL (mg/dL)	99 ± 29	109 ± 34	0.29
HDL (mg/dL)	31 ± 7	42 ± 14	0.002
CRP (mg/dL)	1.10 (0.57-1.55)	0.80 (0.32-1.72)	0.58

Dx: Dialysis, **BMI:** Body mass index, **DMS:** Dialysis malnutrition score, **MAC:** Mid-arm circumference, **URR:** Urea reduction rate, **LDL:** Low-density lipoprotein, **HDL:** High-density lipoprotein, **CRP:** C-reactive protein.

Table III: Comparison of groups with no malnutrition-to-mild malnutrition (DMS: 7-13) and moderate-to-severe malnutrition (DMS: 14-35).

DMS Groups	No change- mild malnutrition (n=22)	Moderate-severe malnutrition (n=18)	p value
Age (years)	51 ± 16	65 ± 17	0.009
Dx vintage (months)	81 (21-117)	93 (22-216)	0.35
BMI (kg/m ²)	26 ± 5	21 ± 4	0.008
Handgrip strength (kg)	28.18 ± 10.21	15.77 ± 8.77	<0.001
MAC (cm)	28.50 ± 4.09	24.61 ± 3.83	0.004
Lean body ratio (%)	38 ± 10	34 ± 8	0.22
Fat tissue ratio (%)	28 ± 14	27 ± 12	0.88
Kt/V	1.44 ± 0.40	1.82 ± 0.48	0.01
URR (%)	71 ± 11	78 ± 8	0.02
Urea (Pre-dx) (mg/dL)	135 ± 38	126 ± 41	0.50
Urea (Post-dx) (mg/dL)	39 ± 17	27 ± 15	0.02
Creatinine (Pre-dx) (mg/dL)	9.90 ± 3.36	7.84 ± 1.43	0.01
Creatinine (Post-dx) (mg/dL)	3.85 ± 1.50	2.31 ± 0.93	0.001
Lymphocyte count (/mm ³)	1677 ± 402	1565 ± 579	0.49
Hemoglobin (g/dL)	11.05 ± 1.30	11.18 ± 1.47	0.77
Uric acid (mg/dL)	6.45 ± 1.38	5.28 ± 1.12	0.006
Albumin (g/dL)	3.61 ± 0.34	3.52 ± 0.30	0.41
T cholesterol (mg/dL)	158 ± 35	181 ± 42	0.06
Triglyceride (mg/dL)	143 ± 71	145 ± 57	0.95
CRP (mg/dL)	0.75 (0.35-1.50)	1.15 (0.55-1.82)	0.18

DMS: Dialysis malnutrition score, **Dx:** Dialysis, **BMI:** Body mass index, **MAC:** Mid-arm circumference, **URR:** Urea reduction rate, **LDL:** Low-density lipoprotein, **HDL:** High-density lipoprotein, **CRP:** C-reactive protein.




Figure 2: Post-dialysis serum creatinine levels were significantly associated with LBR ($r=0.56, p=0.002$).




Figure 3: Post-dialysis serum creatinine levels were significantly associated with HGS ($r=0.73, p<0.001$).

Table IV: Correlation analysis between DMS and various anthropometric and biochemical parameters.

Parameters	Correlation coefficient (r)	p value
Age	0.43	0.005
Dx vintage	0.20	0.24
BMI	-0.28	0.08
Handgrip strength	-0.63	<0.001
MAC	-0.48	0.002
Lean body ratio (%)	-0.31	0.09
Fat tissue ratio (%)	0.15	0.39
Kt/V	0.39	0.01
URR	0.37	0.01
Urea (Pre-dx)	-0.07	0.62
Urea (Post-dx)	-0.37	0.01
Creatinine (Pre-dx)	-0.39	0.01
Creatinine (Post-dx)	-0.57	<0.001
Uric acid	-0.54	<0.001
Total protein	-0.36	0.02
Albumin	-0.24	0.12
Total cholesterol	0.18	0.26
Triglyceride	0.03	0.85
CRP	0.22	0.16

Dx: Dialysis, **BMI:** Body mass index, **MAC:** Mid-arm circumference, **URR:** Urea reduction rate, **CRP:** C-reactive protein.

DISCUSSION

The DMS (SGA) is a practical test of clinical nutritional assessment which is advised for the evaluation of the malnutrition in dialysis patients by the Kidney Disease Outcomes Quality Initiative (K/DOQI) guidelines (9,10). According to DMS, in our study, 22 patients had mild malnutrition and 18 patients had moderate to severe malnutrition. Patients with severe malnutrition were older and they were found to have lower BMI, HGS, MAC and serum uric acid levels. Pre and post-dialysis creatinine levels and only post-dialysis urea levels were lower in the group with severe malnutrition. Interestingly, post-dialysis creatinine and urea levels seemed to be more related to parameters of malnutrition including lean body mass defined by LBR and muscle functions depicted by HGS. Actually, serum creatinine levels are considered to be unreliable as an indicator of muscle mass because they are dialysis dependent. However, in several studies, serum creatinine levels were found to be associated with lean body mass, confirming our




Figure 4: Handgrip strength was significantly negatively correlated with DMS ($r = -0.63, p < 0.001$).




Figure 5: DMS was significantly negatively associated with serum uric acid levels ($r = -0.54, p < 0.001$).

findings (3,11,12). In the univariate correlation analysis, DMS was found to be strongly associated with both anthropometric measurements such as MAC and functions of the muscle mass defined by HGS. However in multivariate analysis, only HGS was found to be associated with DMS.

Table V: Linear regression analysis (Adjusted R² = 0.58; p<0.001) to determine factors independently predicting DMS.

Variables	β	Standardized β	%95 CI		p values
			Lower bound	Upper bound	
Age	0.06	0.22	-0.01	0.13	0.10
Gender	0.92	0.10	-2.26	4.10	0.55
Handgrip strength	-0.19	-0.52	-0.35	-0.03	0.02
MAC	-0.19	-0.19	-0.56	0.18	0.30
LBR	0.02	0.05	-0.15	0.20	0.75
Total protein	-1.41	-0.18	-3.46	0.63	0.16
Uric acid	-0.93	-0.32	-1.77	-0.09	0.03
Kt/V	-1.55	-0.17	-5	1.89	0.35

MAC: Mid-arm circumference, LBR: Lean body ratio.

Table VI: Correlation analysis between handgrip strength and various anthropometric and biochemical parameters.

Parameters	Correlation coefficient (r)	p value
Age	-0.32	0.04
Dx vintage	-0.05	0.74
BMI	0.16	0.32
DMS	-0.63	<0.001
MAC	0.48	0.002
LBR (%)	0.53	0.002
FTR (%)	-0.35	0.04
Kt/V	-0.54	<0.001
URR	-0.56	<0.001
Urea (pre-dx)	0.16	0.31
Urea (post-dx)	0.51	0.001
Creatinine (pre-dx)	0.44	0.005
Creatinine (post-dx)	0.73	<0.001
Uric acid	0.44	0.004
Total protein	0.20	0.21
Albumin	0.05	0.75
Phosphorus	0.30	0.05
Total cholesterol	-0.28	0.07
Triglyceride	0.06	0.69
HDL	-0.33	0.03
LDL	-0.21	0.17
CRP	0.17	0.29
Hemoglobin	0.26	0.10
Ferritin	-0.38	0.01

Dx: Dialysis, BMI: Body mass index, MAC: Mid-arm circumference, LBR: Lean body ratio, FTR: Fat tissue ratio, URR: Urea reduction rate, HDL: High density lipoprotein, LDL: Low density lipoprotein, CRP: C-reactive protein




Figure 6: Handgrip strength was significantly associated with lean body ratio (r = 0.53, p=0.002).

HGS is a simple bedside test that reliably indicates the nutritional status of various patient groups including CKD and dialysis patients (5-7). HGS is known to decrease in patients with CKD. Uremic toxins such as β₂-microglobulin, deficiency of carnitine, disturbances of electrolyte metabolisms, hemodynamic factors related to dialysis procedure, increased inflammation and acidosis may all cause lower values of HGS in HD patients (5,7). Low values of HGS were found to predict mortality in both non-uremic and uremic patients (15,16). Furthermore HGS was also found to predict renal outcomes in non-dialysis CKD patients (17).

Table VII: Linear regression analysis (Adjusted R² =0.71; p<0.001) to determine factors independently predicting handgrip strength.

Variables	β	Standardized β	%95 CI		P values
			Lower bound	Upper bound	
Age	-0.005	-0.006	-0.18	0.17	0.95
Gender	7.93	0.33	0.56	15.29	0.03
DMS	-0.91	-0.33	-1.80	-0.02	0.04
MAC	1.51	0.56	0.48	2.54	0.006
LBR	-0.12	-0.08	-0.81	0.57	0.72
FTR	-0.39	-0.41	-0.88	0.09	0.10
Kt/V	0.75	0.03	-9.91	8.41	0.86
Uric acid	-0.75	-0.09	-3.00	1.50	0.49
HDL	0.20	0.21	-0.06	0.46	0.12
Ferritin	-0.005	-0.12	-0.01	0.005	0.31

DMS: Dialysis malnutrition score, **MAC:** Mid-arm circumference, **LBR:** Lean body ratio, **FTR:** Fat tissue ratio, **HDL:** High density lipoprotein).

In our study, HGS was significantly lower in female patients compared to males as expected and it was found to be significantly associated with MAC, LBR, serum uric acid and post-dialysis serum urea levels. HGS was negatively correlated with age and FTR. In multivariate analysis, the gender, DMS and MAC were found to be significant independent predictors of HGS. Confirming our findings, HGS was found to be negatively associated with DMS scores in several studies (17-19). In several other studies, HGS was directly correlated to LBM in CKD patients in parallel to our findings (15,19-21).

Furthermore, we also found that HGS was negatively associated with FTR but no relationship was present between HGS and BMI. Confirming our findings, in the study by Leal et al., HGS was negatively correlated with body fat ratio and it was not associated with BMI (22). This issue may be explained by the fact that BMI is not a reliable indicator of body composition, in other words it cannot differentiate muscle mass from hypervolemia or fat mass (23). Patients may have normal or high BMI but at the same time they may have decreasing muscle mass and be suffering from PEW. In our study, the negative relationship between HGS and FTR in univariate analysis disappeared in multivariate analysis suggesting a gender-related effect.

Recently, iron accumulation in HD patients (mostly iatrogenic) has been reported to have a negative impact on muscle strength. In the study by Nakagawa et al., serum ferritin levels were found to be independently negatively associated with HGS (24). In a study performed on HD patients with severe proximal myopathy, muscle biopsies showed iron deposition in macrophages and muscle fibers (25). It is hypothesized that

accumulated iron in myoglobin causes free radical formation in muscle leading to impaired muscle functions. Similarly we also found a significant negative association between serum ferritin levels and HGS in univariate analysis. However this effect disappeared in multivariate analysis. Iatrogenic iron overload should be avoided to prevent the possible harmful effects of iron on muscles in HD patients.

In this study, we found that serum uric acid levels were negatively associated with DMS. In multivariate analysis, serum uric acid levels were still significantly predicting DMS independent of Kt/V values. Similarly in the study by Beberashvili et al., serum uric acid levels were negatively correlated with the malnutrition-inflammation score (26). In our study, serum uric acid levels were also correlated with HGS in univariate analysis but this correlation disappeared in multivariate analysis. In the study by Beberashvili et al., serum uric acid levels were found to be positively associated with body composition parameters such as BMI, MAC, fat mass, lean body mass and HGS (26). Uric acid is the end product of purine metabolism. Increased serum uric acid levels are known to be associated with an increased risk for CKD (27,28), and cardiovascular disease in the general population (29,30). However, in HD patients, low uric acid levels have been found to be associated with higher risk of mortality (31-33). Higher serum uric acid levels have been suggested to reflect a better nutritional status in HD patients.

Adequacy of dialysis, defined by Kt/V and URR, was found to be higher in patients with more severe malnutrition determined by DMS. Kt/V and URR were negatively associated with HGS in univariate analysis. However, parameters of dialysis adequacy did not independently predict HGS in multivariate regression

analysis. Confirming our findings, single pool Kt/V was found to be negatively correlated with both skeletal muscle index and HGS in the study by Morishita et al. (34). The sole determinant of Kt/V is not only the dialysis efficacy and the muscle mass of the patients also affects the Kt/V values. HD patients with high muscle mass have been reported to have low Kt/V regardless of dialysis efficacy (35,36). There are also controversies in terms of the relationship between Kt/V and HGS; Qureshi et al. (6) could not find an association between HGS and Kt/V.

The observational and cross-sectional design of the study and a relatively low sample size may be considered as the limitations of this study. Further studies with long-term follow-up data will contribute to our knowledge of malnutrition in HD patients.

In conclusion, HGS and serum uric acid levels were significantly associated with dialysis malnutrition scores. HGS is an easy and reliable test for the evaluation of malnutrition in HD patients. BMI may be misleading as a parameter of malnutrition in HD patients and body composition analysis with BIA is therefore crucial for follow-up of LBR. Evaluation of the effects of dialysis adequacy on malnutrition by the use of Kt/V may be difficult and misleading.

REFERENCES

1. Fouque D, Kalantar-Zadeh K, Kopple J, Cano N, Chauveau P, Cuppari L, Franch H, Guarnieri G, Ikizler TA, Kaysen G, Lindholm B, Massy Z, Mitch W, Pineda E, Stenvinkel P, Treviño-Becerra A, Wanner C: A proposed nomenclature and diagnostic criteria for protein-energy wasting in acute and chronic kidney disease. *Kidney Int* 2008;73:391-398
2. Kalantar-Zadeh K, Ikizler TA, Block G, Avram MM, Kopple JD: Malnutrition-inflammation complex syndrome in dialysis patients: Causes and consequences. *Am J Kidney Dis* 2003;42:864-881
3. Rymarz A, Bartoszewicz Z, Szamotulska K, Niemczyk S: The associations between body cell mass and nutritional and inflammatory markers in patients with chronic kidney disease and in subjects without kidney disease. *J Ren Nutr* 2016;26:87-92
4. Bohannon RW: Dynamometer measurements of hand-grip strength predict multiple outcomes. *Percept Mot Skills* 2001;93:323-328
5. Wang AY, Sea MM, Ho ZS, Lui SF, Li PK, Woo J: Evaluation of handgrip strength as a nutritional marker and prognostic indicator in peritoneal dialysis patients. *Am J Clin Nutr* 2005;81:79-86
6. Qureshi AR, Alvestrand A, Danielsson A, Divino-Filho JC, Gutierrez A, Lindholm B, Bergström J: Factors predicting malnutrition in hemodialysis patients: A cross-sectional study. *Kidney Int* 1998;53:773-782
7. Leal VO, Mafra D, Fouque D, Anjos LA: Use of handgrip strength in the assessment of the muscle function of chronic kidney disease patients on dialysis: A systematic review. *Nephrol Dial Transplant*. 2011;26:1354-1360
8. Kalantar-Zadeh K, Kleiner M, Dunne E, Lee GH, Luft FC: A modified quantitative subjective global assessment of nutrition for dialysis patients. *Nephrol Dial Transplant* 1999;14:1732-1738
9. Riella MC: Nutritional evaluation of patients receiving dialysis for the management of protein-energy wasting: What is old and what is new? *J Ren Nutr* 2013;23:195-198
10. K/DOQI, National Kidney Foundation: Clinical practice guidelines for nutrition in chronic renal failure. *Am J Kidney Dis* 2000;35:S1-S140
11. Noori N, Kovesdy CP, Dukkipati R, Kim Y, Duong U, Bross R, Oreopoulos A, Luna A, Benner D, Kopple JD, Kalantar-Zadeh K: Survival predictability of lean and fat mass in men and women undergoing maintenance hemodialysis. *Am J Clin Nutr* 2010;92:1060-1070
12. Patel SS, Molnar MZ, Tayek JA, Ix JH, Noori N, Benner D, Heymsfield S, Kopple JD, Kovesdy CP, Kalantar-Zadeh K: Serum creatinine as a marker of muscle mass in chronic kidney disease: Results of a cross-sectional study and review of literature. *J Cachexia Sarcopenia Muscle* 2013;4:19-29
13. Limaye V, Frankham A, Disney A, Pile K: Evaluation of hand function in patients undergoing long term haemodialysis. *Ann Rheum Dis* 2001;60:278-280
14. Constatin-Teodosiu D, Young S, Wellock F, Short AH, Burden RP, Morgan AG, Greenhaff PL: Gender and age differences in plasma carnitine, muscle strength, and exercise tolerance in haemodialysis patients. *Nephrol Dial Transplant* 2002;17:1808-1813
15. Heimbürger O, Qureshi AR, Blauer WS, Berglund L, Stenvinkel P: Hand-grip muscle strength, lean body mass, and plasma proteins as markers of nutritional status in patients with chronic renal failure close to start of dialysis therapy. *Am J Kidney Dis* 2000;36:1213-1225
16. Vogt BP, Borges MC, Goés CR, Caramori JC: Handgrip strength is an independent predictor of all-cause mortality in maintenance dialysis patients. *Clin Nutr* 2016;35:1429-1433
17. Chang YT, Wu HL, Guo HR, Cheng YY, Tseng CC, Wang MC, Lin CY, Sung JM: Handgrip strength is an independent predictor of renal outcomes in patients with chronic kidney diseases. *Nephrol Dial Transplant* 2011;26:3588-3595
18. Pagels A, Heiwe S, Hylander B: Nutritional status of pre-dialysis patients. *J Ren Care* 2006;32:162-166
19. Stenvinkel P, Barany P, Chung SH, Lindholm B, Heimbürger O: A comparative analysis of nutritional parameters as predictor of outcome in male and female ESRD patients. *Nephrol Dial Transplant* 2002;17:1266-1274
20. Konings CJ, Kooman JP, Schonck M, van Kreef B, Heidendal GA, Cheriex EC, van der Sande FM, Leunissen KM: Influence of fluid status on techniques used to assess body composition in peritoneal dialysis patients. *Perit Dial Int* 2003;23:184-190
21. Jones CH, Newstead CG, Will EJ, Smye SW, Davison AM: Assessment of nutritional status in CAPD patients: Serum albumin is not a useful measure. *Nephrol Dial Transplant* 1997;12:1406-1413

22. Leal VO, Stockler-Pinto MB, Farage NE, Aranha LN, Fouque D, Anjos LA, Mafrá D: Handgrip strength and its dialysis determinants in hemodialysis patients. *Nutrition* 2011;27:1125-129
23. Mafrá D, Guebre-Egziabher F, Fouque D: Body mass index, muscle and fat in chronic kidney disease: Questions about survival. *Nephrol Dial Trans* 2008;23:2461-2466
24. Nakagawa C, Inaba M, Ishimura E, Yamakawa T, Shoji S, Okuno S: Association of increased serum ferritin with impaired muscle strength/quality in hemodialysis patients. *J Ren Nutr* 2016;26:253-257
25. Bregman H, Gelfand MC, Winchester JF, Manz HJ, Kneppshield JH, Schreiner GE: Iron-overload-associated myopathy in patients on maintenance haemodialysis: A histocompatibility-linked disorder. *Lancet* 1980;2:882-885
26. Beberashvili I, Sinuani I, Azar A, Shapiro G, Feldman L, Stav K, Sandbank J, Averbukh Z: Serum uric acid as a clinically useful nutritional marker and predictor of outcome in maintenance hemodialysis patients. *Nutrition* 2015;31:138-147
27. Feig DI: Uric acid: A novel mediator and marker of risk in chronic kidney disease? *Curr Opin Nephrol Hypertens* 2009;18:526-530
28. Kanbay M, Ozkara A, Selcoki Y, Isik B, Turgut F, Bavbek N, Uz E, Akcay A, Yigitoglu R, Covic A: Effect of treatment of hyperuricemia with allopurinol on blood pressure, creatinine clearance, and proteinuria in patients with normal renal functions. *Int Urol Nephrol* 2007;39:1227-1233
29. Feig DI, Kang DH, Johnson RJ: Uric acid and cardiovascular risk. *N Engl J Med* 2008;359:1811-1821
30. Kanbay M1, Yilmaz MI, Sonmez A, Turgut F, Saglam M, Cakir E, Yenicesu M, Covic A, Jalal D, Johnson RJ: Serum uric acid level and endothelial dysfunction in patients with nondiabetic chronic kidney disease. *Am J Nephrol* 2011;33:298-304
31. Suliman ME, Johnson RJ, García-López E, Qureshi AR, Molinaei H, Carrero JJ, Heimbürger O, Bárány P, Axelsson J, Lindholm B, Stenvinkel P: J-shaped mortality relationship for uric acid in CKD. *Am J Kidney Dis* 2006;48:761-771
32. Latif W, Karaboyas A, Tong L, Winchester JF, Arrington CJ, Pisoni RL, Marshall MR, Kleophas W, Levin NW, Sen A, Robinson BM, Saran R: Uric acid levels and all-cause and cardiovascular mortality in the hemodialysis population. *Clin J Am Soc Nephrol* 2011;6:2470-2477
33. Lee SM, Lee AL, Winters TJ, Tam E, Jaleel M, Stenvinkel P, Johnson RJ: Low serum uric acid level is a risk factor for death in incident hemodialysis patients. *Am J Nephrol* 2009;29:79-85
34. Morishita Y, Kubo K, Haga Y, Miki A, Ishibashi K, Kusano E, Nagata D. Skeletal muscle loss is negatively associated with single-pool Kt/V and dialysis duration in hemodialysis patients. *Ther Apher Dial* 2014;18:612-7
35. Chertow GM, Owen WF, Lazarus JM, Lew NL, Lowrie EG. Exploring the reverse J-shaped curve between urea reduction ratio and mortality. *Kidney Int* 1999;56:1872-8
36. Li Z, Lew NL, Lazarus JM, Lowrie EG. Comparing the urea reduction ratio and the urea product as outcome-based measures of hemodialysis dose. *Am J Kidney Dis* 2000;35:598-605